

COVID-19 Scenario 1

Cohort Ward PPE Donning Instructions

Please follow PPE donning procedure as instructed to be undertaken in pairs. Perform hand hygiene before undertaking the following steps:

Before you enter the ward

Before seeing a COVID-19 patient

Step 4.

Put on surgical cap over the straps of the mask and eye protection.

4

Step 5.

Put on apron. Break neck strap and tie high up to cover upper chest.

Step 3.

Put on eye protection.

3

Step 2.

Put on fluid resistant surgical face mask.

2

Step 1.

Please ensure you are wearing scrubs.

1

Step 6.

Put on gloves.

6

COVID-19 Scenario 1

Cohort Ward PPE Doffing Instructions

Please follow PPE doffing procedure. Read instructions out loud with a buddy while removing PPE.
PERFORM HAND HYGIENE AFTER EACH STEP OF THE DOFFING PROCESS. Scrubs and PPE should not be worn outside of this clinical area.

Step 3: outside room

Remove surgical cap. Perform hand hygiene.



Step 2: inside room

Remove apron. Perform hand hygiene.



Step 4: outside room

Remove eye protection. The outside of the face shield is contaminated and should not be touched. Perform hand hygiene. Clean eye protection with green Clinell wipe for **60 seconds**. Once dry can be re-used. Perform hand hygiene.




Step 1: inside room

Remove gloves. Perform hand hygiene.

Grasp the outside of glove with the opposite gloved hand and peel off.

Hold the removed glove in the remaining gloved hand. Slide the fingers of the un-gloved hand under the remaining glove at the wrist and peel off. Discard as clinical waste. Perform hand hygiene.



Step 5: outside room

Only touch straps when removing mask. Perform hand hygiene.

